Climate Change Projections for Individual Tree Species

Missouri-Nishnabotna Watershed in Iowa



This region's forests will be affected by a changing climate and other stressors during this century. The National Climate Assessment describes how a changing climate can increase the vulnerability of forests in the Midwest (Wilson et al. 2023: doi.org/10.7930/NCA5.2023.CH24). This report includes information on the current landscape, observed climate trends, and a range of projected future climates. This handout summarizes data from the U.S. Forest Service's Climate Change Tree Atlas (doi.org/10.2737/Climate-Change-Tree-Atlas-v4). Two climate

scenarios are presented to "bracket" a range of possible futures. These future climate projections (2070 to 2099) provide information about how individual tree species may respond to a changing climate. Results for "low" and "high" emissions scenarios can be compared on the reverse side of this handout.

The Tree Atlas provides information to interpret tree species changes:

- **SUITABLE HABITAT** calculated based on 45 variables that explain where conditions exist for a species, including soils, landforms, and climate variables.
- ADAPTABILITY based on life-history traits that might increase or decrease tolerance of expected changes, such as the ability to withstand different forms of disturbance
- CAPABILITY a rating of the species' ability to cope or persist with climate change in this region based on suitable habitat change (statistical modeling), adaptability (literature review and expert opinion), and abundance (inventory data). The capability rating is modified by abundance information; ratings are downgraded for rare species and upgraded for abundant species. Capability rating may not appropriately reflect the outlook for species with emerging severe forest health issues, such as ash species affected by emerald ash borer. See the table to the right for ratings.

CLIMATE CHANGE CAPABILITY TABLE.

Capability is a rating of the species' ability to cope or persist with climate change. Species are organized into poor, fair, good, and mixed capability ratings. Species with new suitable habitat or low model reliability are excluded from this table. See the Tree Species Projections table legend on the following page for more information on ratings.

NOTE: Capability rating may not reflect severe forest health issues (e.g., emerald ash borer).

POOR CAPABILITY							
American basswood	Red pine						
Black cherry	Shagbark hickory						
Chinkapin oak	Shingle oak						
Flowering dogwood	White ash						
Northern pin oak							
FAIR CAPABILITY							
American elm	Eastern redcedar						
Black oak	White oak						
GOOD CAPABILITY							
Blackjack oak	Post oak						
Hackberry	Sugarberry						
Osage-orange							
MIXED CAPABILITY							
Bur oak	Sugar maple						
Northern red oak							

• **MIGRATION POTENTIAL MODEL** - when combined with habitat suitability, an estimate of a species' colonization likelihood for new habitats. This rating can be helpful for assisted migration or focused management.

Remember that models are just tools, and they're not perfect. Model projections can't account for all factors that influence future species success. If a species is rare or confined to a small area, model results may be less reliable. These factors, and others, could cause a particular species to perform better or worse than a model projects. Human choices will also continue to influence forest distribution, especially for tree species that are projected to increase. Planting programs may assist the movement of future-adapted species, but this will depend on management decisions. Despite these limits, models provide useful information about future expectations. It's perhaps best to think of these projections as indicators of possibility and potential change.

CREDIT: This handout summarizes the full model results for the Missouri-Nishnabotna Watershed in Iowa. Data provided by the USDA Forest Service (M.P. Peters, A.M. Prasad, S.N. Matthews, & L.R. Iverson) as part of the Climate Change Tree Atlas (doi.org/10.2737/Climate-Change-Tree-Atlas-v4). Models and variables are described in Iverson et al. 2019 and Peters et al. 2019 (available at <u>fs.usda.gov/nrs/atlas/products/pubs</u>). More information on vulnerability and adaptation in the region can be found at <u>forestadaptation.org/northwoods</u>.







Tree Species Projections Table

Information presented in the table is from the Climate Change Tree Atlas regional summaries, more details at fs.usda.gov/nrs/atlas/combined/resources/summaries.

ADAPTABILITY: Life-history factors, such as the ability to respond favorably to disturbance, that are not included in the Tree Atlas model and may make a species more or less able to adapt to future stressors.

- + **HIGH** Species may perform better than modeled
- **MEDIUM**
- **LOW** Species may perform worse than modeled

HABITAT CHANGE: Projected change in suitable habitat between current and potential future conditions.

- ▲ INCREASE Projected increase of >20% by 2100
- **DECREASE** Projected decrease of >20% by 2100
- NO CHANGE Projected change of <20% by 2100
- **NEW HABITAT** Tree Atlas projects new habitat for species not currently present

ABUNDANCE: Based on Forest Inventory Analysis (FIA) summed Importance Value data, calibrated to a standard geographic area.

- + ABUNDANT
- COMMON
- **RARE**

CAPABILITY: An overall rating that describes a species' ability to cope or persist with climate change based on suitable habitat change class, adaptability, and abundance within this region. Capability may not reflect severe forest health issues.

- △ GOOD Increasing suitable habitat, medium or high adaptability, and common or abundant
- O FAIR Mixed combinations, such as a rare species with increasing suitable habitat and medium adaptability
- **▼ POOR** Decreasing suitable habitat, medium or low adaptability, and uncommon or rare

			LOW CLIMATE CHANGE (RCP 4.5)		HIGH CLIMATE CHANGE (RCP 8.5)					LOW CLIMATE CHANGE (RCP 4.5)		HIGH CLIMATE CHANGE (RCP 8.5)	
SPECIES	ADAPT	ABUN	HABITAT CHANGE		HABITAT CHANGE	CAPABILITY	SPECIES	ADAPT	ABUN	HABITAT CHANGE		HABITAT CHANGE	CAPABILITY
American basswood	•	_	_	$\overline{\nabla}$	V	∇	Northern pin oak	+	_	_	∇	V	∇
American elm	•	•	•	0	•	0	Northern red oak	+	_	•	0	_	0
Ashe juniper	•	_	*		*		Ohio buckeye	•	_	_	∇	_	∇
Bitternut hickory*	+	_	•	0	•	0	Osage-orange	+	_	A	Δ	_	Δ
Black cherry	-	-	_	∇	_	∇	Pecan*	_	_	*		*	
Black hickory	•	_	*		*		Pignut hickory	•	_	*		*	
Black locust*	•	_	•	∇	•	∇	Pin oak*	_	_	_	∇	•	0
Black oak	•	_	_	0	_	0	Post oak	+	_		Δ		Δ
Black walnut*	•		•	0	•	0	Red mulberry*	•	_	•	∇		∇
Black willow*	_	_	•	∇		$\overline{\nabla}$	Red pine	_	_	•	∇	_	∇
Blackjack oak	+	_	<u> </u>	Δ	•	Δ	Sassafras*	•	_	*		*	
Boxelder*	+	_	_	∇	•	0	Serviceberry*	•	_	_	∇	_	∇
Bur oak	+	_	<u> </u>	Δ	•	0	Shagbark hickory	•	_	_	∇		∇
Cedar elm	_	_	*		*		Shingle oak	•	_	•	∇	_	∇
Chinkapin oak	•	_	_	∇	_	$\overline{\nabla}$	Shumard oak*	+	_	*		*	
Common persimmon*	+	_	<u> </u>	Δ		Δ	Silver maple*	+	_	•	0	•	0
Eastern cottonwood*		_	•	∇	•	$\overline{\nabla}$	Slippery elm*	•	_	•	∇		0
Eastern redbud*	•	_	•	∇	•	$\overline{\nabla}$	Sugar maple	+	_	•	0		Δ
Eastern redcedar	•	_	<u> </u>	0		0	Sugarberry	•	_	*		*	
Flowering dogwood	•	_	_	∇	_	$\overline{\nabla}$	Swamp white oak*	•	_	_	∇	_	∇
Green ash*		_	<u> </u>	Δ		Δ	Sycamore*	•	_	•	∇	•	∇
Hackberry	+	•	•	Δ	•	Δ	Water oak	+	_	*		*	
Honeylocust*	+	•	_	0	_	0	White ash	_	_	•	∇	•	∇
Ironwood*	+	_	_	∇	_	∇	White oak	+	_	•	0	•	0
Mockernut hickory†	+	_	*		*		Winged elm	•	_	*		*	

^{*}Species with low model reliability based on five statistical metrics of the habitat models that affect change class.

[†]Species is likely present but not currently included in the FIA database.